

Lesson 4.2 - Coterminal angles

Learning Objectives: SWBAT

1. Determine and sketch angles that are coterminal to a given angle
2. Determine complimentary and supplementary angles (in radians)

An **angle** is determined by rotating a ray (half-line) about its endpoint. The starting position of the ray is the **initial side** of the angle, and the position after rotation is the **terminal side**, as shown in Figure 4.1. The endpoint of the ray is the **vertex** of the angle. This perception of an angle fits a coordinate system in which the origin is the vertex and the initial side coincides with the positive x -axis. Such an angle is in **standard position**, as shown in Figure 4.2. **Positive angles** are generated by counterclockwise rotation, and **negative angles** by clockwise rotation, as shown in Figure 4.3. Angles are labeled with Greek letters such as α (alpha), β (beta), and θ (theta), as well as uppercase letters such as A , B , and C . In Figure 4.4, note that angles α and β have the same initial and terminal sides. Such angles are **coterminal**.




Figure 4.1




Figure 4.2




Figure 4.3






Figure 4.4

Two positive angles α and β are **complementary** (complements of each other) if their sum is $\pi/2$. Two positive angles are **supplementary** (supplements of each other) if their sum is π . See Figure 4.12.



Complementary angles

Figure 4.12



Supplementary angles

Lesson 4.2 - Coterminal angles

Examples: How to find coterminal angle(s) of a given angle

- a. For the positive angle $\theta = \frac{13\pi}{6}$, subtract 2π to obtain a coterminal angle

$$\frac{13\pi}{6} - 2\pi = \frac{\pi}{6}$$

See Figure 4.9.




Figure 4.9

- b. For the positive angle $\theta = \frac{3\pi}{4}$, subtract 2π to obtain a coterminal angle

$$\frac{3\pi}{4} - 2\pi = -\frac{5\pi}{4}$$

See Figure 4.10.




Figure 4.10

- c. For the negative angle $\theta = -\frac{2\pi}{3}$, add 2π to obtain a coterminal angle

$$-\frac{2\pi}{3} + 2\pi = \frac{4\pi}{3}$$

See Figure 4.11.






Figure 4.11

Your Turn: For the negative angle $-\frac{5\pi}{4}$, determine two coterminal angles

there are many correct answers



$$-\frac{5\pi}{4} + 2\pi = \frac{3\pi}{4}$$

$$-\frac{5\pi}{4} + 2\pi = \frac{-13\pi}{4}$$

Lesson 4.2 - Coterminal angles

Examples: How to find complimentary and supplementary angles (in radians)

If possible, find the complement and the supplement of (a) $\frac{2\pi}{5}$ and (b) $\frac{4\pi}{5}$.

Solution

a. The complement of $\frac{2\pi}{5}$ is

$$\begin{aligned} \frac{\pi}{2} - \frac{2\pi}{5} &= \frac{5\pi}{10} - \frac{4\pi}{10} \\ &= \frac{\pi}{10} \end{aligned}$$

The supplement of $\frac{2\pi}{5}$ is

$$\begin{aligned} \pi - \frac{2\pi}{5} &= \frac{5\pi}{5} - \frac{2\pi}{5} \\ &= \frac{3\pi}{5} \end{aligned}$$



b. Because $4\pi/5$ is greater than $\pi/2$, it has no complement. (Remember that complements are *positive* angles.) The supplement is

$$\begin{aligned} \pi - \frac{4\pi}{5} &= \frac{5\pi}{5} - \frac{4\pi}{5} \\ &= \frac{\pi}{5} \end{aligned}$$

Practice:



In Exercises 11–14, determine two coterminal angles in radian measure (one positive and one negative) for each angle. (There are many correct answers).

$$\begin{aligned} \frac{11\pi}{6} + 2\pi &= \frac{13\pi}{6} \\ \frac{11\pi}{6} - 2\pi &= -\frac{11\pi}{6} \end{aligned}$$



$$\begin{aligned} \frac{2\pi}{3} + 2\pi &= \frac{8\pi}{3} \\ \frac{2\pi}{3} - 2\pi &= -\frac{4\pi}{3} \end{aligned}$$

$$\begin{aligned} \frac{7\pi}{6} + 2\pi &= \frac{19\pi}{6} \\ \frac{7\pi}{6} - 2\pi &= -\frac{3\pi}{4} \end{aligned}$$



$$\begin{aligned} \frac{5\pi}{4} + 2\pi &= \frac{13\pi}{4} \\ \frac{5\pi}{4} - 2\pi &= -\frac{3\pi}{4} \end{aligned}$$

$$\begin{aligned} -\frac{9\pi}{4} + 2\pi &= -\frac{\pi}{4} \\ -\frac{9\pi}{4} - 2\pi &= \frac{7\pi}{4} \end{aligned}$$

13. (a) $-\frac{9\pi}{4}$

(b) $-\frac{2\pi}{15}$

$$-\frac{2\pi}{15} + 2\pi = \frac{28\pi}{15}$$

$$-\frac{2\pi}{15} - 2\pi = -\frac{32\pi}{15}$$

14. (a) $\frac{7\pi}{8}$

(b) $\frac{8\pi}{45}$

$$\frac{8\pi}{45} + 2\pi = \frac{98\pi}{45}$$

$$\frac{8\pi}{45} - 2\pi = -\frac{82\pi}{45}$$

$$\frac{7\pi}{8} + 2\pi = \frac{23\pi}{8}$$

$$\frac{7\pi}{8} - 2\pi = -\frac{9\pi}{8}$$

Lesson 4.2 - Coterminal angles

Practice:

In Exercises 15–20, find (if possible) the complement and supplement of the angle.

15. $\frac{\pi}{3}$

Comp: $\frac{\pi}{2} - \frac{\pi}{3} = \frac{\pi}{6}$

Supp: $\pi - \frac{\pi}{3} = \frac{2\pi}{3}$

16. $\frac{3\pi}{4}$

Comp: ~~$\frac{\pi}{2} - \frac{3\pi}{4} = \text{Not possible}$~~ $\frac{\pi}{2}$

Supp: $\pi - \frac{3\pi}{4} = \frac{\pi}{4}$

17. $\frac{\pi}{6}$

Comp: $\frac{\pi}{2} - \frac{\pi}{6} = \frac{\pi}{3}$

Supp: $\pi - \frac{\pi}{6} = \frac{5\pi}{6}$


18. $\frac{2\pi}{3}$

Comp: Not possible

Supp: $\pi - \frac{2\pi}{3} = \frac{\pi}{3}$

In Exercises 31–34, determine two coterminal angles in degree measure (one positive and one negative) for each angle. (There are many correct answers).


31. (a)



$52 + 360 = 412^\circ$

$52 - 360 = -308^\circ$


(b)



$-36 + 360 = 324^\circ$

$-36 - 360 = -396^\circ$


32. (a)



$114 + 360 = 474^\circ$

$114 - 360 = -246^\circ$

(b)



$-390 + 360 = -30^\circ$

$-390 - 360 = -750^\circ$

33. (a) 300°

$300 + 360 = 660^\circ$

$300 - 360 = -60^\circ$

(b) 230°

$230 + 360 = 590^\circ$

$230 - 360 = -130^\circ$

34. (a) -445°

$-445 + 360 = -85^\circ$


$-445 - 360 = -805^\circ$

(b) -740°

$-740 + 360 = -380^\circ$

$-740 - 360 = -1100^\circ$

75. Find each angle (in radians) shown on the unit circle.



$30^\circ = \frac{\pi}{6}$

$45^\circ = \frac{\pi}{4}$

$60^\circ = \frac{\pi}{3}$

$90^\circ = \frac{\pi}{2}$

$135^\circ = \frac{3\pi}{4}$

$180^\circ = \pi$

$210^\circ = \frac{7\pi}{6}$

$270^\circ = \frac{3\pi}{2}$

$330^\circ = \frac{11\pi}{6}$