


## GRAPHS OF PARENT FUNCTIONS

### Linear Function

$$f(x) = mx + b$$



Domain:  $(-\infty, \infty)$

Range:  $(-\infty, \infty)$

$x$ -intercept:  $(-b/m, 0)$


$y$ -intercept:  $(0, b)$

Increasing when  $m > 0$

Decreasing when  $m < 0$

### Absolute Value Function

$$f(x) = |x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$$



Domain:  $(-\infty, \infty)$

Range:  $[0, \infty)$

Intercept:  $(0, 0)$

Decreasing on  $(-\infty, 0)$


Increasing on  $(0, \infty)$

Even function

$y$ -axis symmetry

### Square Root Function

$$f(x) = \sqrt{x}$$



Domain:  $[0, \infty)$


Range:  $[0, \infty)$

Intercept:  $(0, 0)$

Increasing on  $(0, \infty)$

### Greatest Integer Function

$$f(x) = \llbracket x \rrbracket$$



Domain:  $(-\infty, \infty)$

Range: the set of integers

$x$ -intercepts: in the interval  $[0, 1)$


$y$ -intercept:  $(0, 0)$

Constant between each pair of consecutive integers

Jumps vertically one unit at each integer value

### Quadratic (Squaring) Function

$$f(x) = ax^2$$



Domain:  $(-\infty, \infty)$

Range ( $a > 0$ ):  $[0, \infty)$

Range ( $a < 0$ ):  $(-\infty, 0]$

Intercept:  $(0, 0)$

Decreasing on  $(-\infty, 0)$  for  $a > 0$

Increasing on  $(0, \infty)$  for  $a > 0$

Increasing on  $(-\infty, 0)$  for  $a < 0$

Decreasing on  $(0, \infty)$  for  $a < 0$


Even function

$y$ -axis symmetry

Relative minimum ( $a > 0$ ), relative maximum ( $a < 0$ ), or vertex:  $(0, 0)$

### Cubic Function

$$f(x) = x^3$$



Domain:  $(-\infty, \infty)$

Range:  $(-\infty, \infty)$

Intercept:  $(0, 0)$

Increasing on  $(-\infty, \infty)$


Odd function

Origin symmetry

## GRAPHS OF PARENT FUNCTIONS

### Rational (Reciprocal) Function


$$f(x) = \frac{1}{x}$$



Domain:  $(-\infty, 0) \cup (0, \infty)$   
 Range:  $(-\infty, 0) \cup (0, \infty)$   
 No intercepts  
 Decreasing on  $(-\infty, 0)$  and  $(0, \infty)$   
 Odd function  
 Origin symmetry  
 Vertical asymptote:  $y$ -axis  
 Horizontal asymptote:  $x$ -axis

### Exponential Function


$$f(x) = a^x, a > 0, a \neq 1$$



Domain:  $(-\infty, \infty)$   
 Range:  $(0, \infty)$   
 Intercept:  $(0, 1)$   
 Increasing on  $(-\infty, \infty)$   
 for  $f(x) = a^x$   
 Decreasing on  $(-\infty, \infty)$   
 for  $f(x) = a^{-x}$   
 Horizontal asymptote:  $x$ -axis  
 Continuous

### Logarithmic Function


$$f(x) = \log_a x, a > 0, a \neq 1$$



Domain:  $(0, \infty)$   
 Range:  $(-\infty, \infty)$   
 Intercept:  $(1, 0)$   
 Increasing on  $(0, \infty)$   
 Vertical asymptote:  $y$ -axis  
 Continuous  
 Reflection of graph of  $f(x) = a^x$   
 in the line  $y = x$

### Sine Function


$$f(x) = \sin x$$



Domain:  $(-\infty, \infty)$   
 Range:  $[-1, 1]$   
 Period:  $2\pi$   
 x-intercepts:  $(n\pi, 0)$   
 y-intercept:  $(0, 0)$   
 Odd function  
 Origin symmetry

### Cosine Function


$$f(x) = \cos x$$



Domain:  $(-\infty, \infty)$   
 Range:  $[-1, 1]$   
 Period:  $2\pi$   
 x-intercepts:  $\left(\frac{\pi}{2} + n\pi, 0\right)$   
 y-intercept:  $(0, 1)$   
 Even function  
 y-axis symmetry

### Tangent Function

$$f(x) = \tan x$$




Domain: all  $x \neq \frac{\pi}{2} + n\pi$   
 Range:  $(-\infty, \infty)$   
 Period:  $\pi$   
 x-intercepts:  $(n\pi, 0)$   
 y-intercept:  $(0, 0)$   
 Vertical asymptotes:  
 $x = \frac{\pi}{2} + n\pi$   
 Odd function  
 Origin symmetry

## GRAPHS OF PARENT FUNCTIONS

### Cosecant Function

$$f(x) = \csc x$$



Domain: all  $x \neq n\pi$

Range:  $(-\infty, -1] \cup [1, \infty)$

Period:  $2\pi$

No intercepts


Vertical asymptotes:  $x = n\pi$

Odd function

Origin symmetry

### Secant Function

$$f(x) = \sec x$$



Domain: all  $x \neq \frac{\pi}{2} + n\pi$

Range:  $(-\infty, -1] \cup [1, \infty)$

Period:  $2\pi$

$y$ -intercept:  $(0, 1)$

Vertical asymptotes:


$$x = \frac{\pi}{2} + n\pi$$

Even function

$y$ -axis symmetry

### Cotangent Function

$$f(x) = \cot x$$



Domain: all  $x \neq n\pi$

Range:  $(-\infty, \infty)$

Period:  $\pi$

$$x\text{-intercepts: } \left(\frac{\pi}{2} + n\pi, 0\right)$$


Vertical asymptotes:  $x = n\pi$

Odd function

Origin symmetry

### Inverse Sine Function

$$f(x) = \arcsin x$$



Domain:  $[-1, 1]$

$$\text{Range: } \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$$


Intercept:  $(0, 0)$

Odd function

Origin symmetry

### Inverse Cosine Function

$$f(x) = \arccos x$$




Domain:  $[-1, 1]$

$$\text{Range: } [0, \pi]$$

$$y\text{-intercept: } \left(0, \frac{\pi}{2}\right)$$

### Inverse Tangent Function

$$f(x) = \arctan x$$



Domain:  $(-\infty, \infty)$

$$\text{Range: } \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$

Intercept:  $(0, 0)$

Horizontal asymptotes:

$$y = \pm \frac{\pi}{2}$$

Odd function

Origin symmetry